

The Szpiro inequality for higher genus fibrations

Arnaud BEAUVILLE

Introduction

The aim of this note is to prove the following result:

Proposition .— *Let $f : S \rightarrow B$ be a non-trivial semi-stable fibration of genus $g \geq 2$, N the number of critical points of f and s the number of singular fibres. Then*

$$N < (4g + 2)(s + 2g(B) - 2) .$$

Recall that a semi-stable fibration of genus g is a surjective holomorphic map of a smooth projective surface S onto a smooth curve B , whose generic fibre is a smooth curve of genus g and whose singular fibres are allowed only ordinary double points; moreover we impose that each smooth rational curve contained in a fibre meets the rest of the fibre in at least 2 points (otherwise by blowing up non-critical points of f in a singular fibre we could arbitrarily increase N keeping s fixed).

The corresponding inequality $N \leq 6(s + 2g(B) - 2)$ in the case $g = 1$ has been observed by Szpiro; it was motivated by the case of curves over a number field, where an analogous inequality would have far-reaching consequences [S]. The higher genus case is considered in the recent preprint [BKP], where the authors prove the slightly weaker inequality $N \leq (4g + 2)s$ for hyperelliptic fibrations over $\mathbf{P}^1$. Their method is topological, and in fact the result applies in the much wider context of symplectic Lefschetz fibrations. We will show that in the more restricted algebraic-geometric set-up, the Proposition is a direct consequence of two classical inequalities in surface theory. It would be interesting to know whether the proof of [BKP] can be extended to non-hyperelliptic fibrations.

Proof

The main numerical invariants of a surface S are the square of the canonical bundle K_S , the Euler-Poincaré characteristic $\chi(\mathcal{O}_S)$ and the topological Euler-Poincaré characteristic $e(S)$; they are linked by the Noether formula $12\chi(\mathcal{O}_S) = K_S^2 + e(S)$. For a semi-stable fibration $f : S \rightarrow B$ it has become customary to modify these invariants as follows. Let b be the genus of B , and $K_f = K_X \otimes f^*K_B^{-1}$ the relative canonical bundle of X over B ; then we consider:

$$\begin{aligned} K_f^2 &= K_X^2 - 8(b - 1)(g - 1) \\ \chi_f &:= \deg f_*(K_f) = \chi(\mathcal{O}_X) - (b - 1)(g - 1) \\ e_f &:= N = e(X) - 4(b - 1)(g - 1) . \end{aligned}$$

Observe that we have again $12\chi_f = K_f^2 + e_f$. We will use the Xiao inequality ([X], Theorem 2)

$$K_f^2 \geq \left(4 - \frac{4}{g}\right) \chi_f$$

and the “strict canonical class inequality” ([T], lemma 3.1)

$$K_f^2 < 2(g-1)(s+2b-2) \quad \text{for } s > 0.$$

Let us prove the Proposition. If $s = 0$, we have $N = 0$ and $g(B) \geq 2$ (otherwise the fibration would be trivial), so the inequality of the Proposition holds. Assume $s > 0$; the Xiao inequality gives

$$3gK_f^2 \geq 12(g-1)\chi_f = (g-1)(K_f^2 + e_f),$$

hence, using the strict canonical class inequality,

$$N = e_f \leq \frac{2g+1}{g-1} K_f^2 < (4g+2)(s+2b-2). \quad \blacksquare$$

Example.— We constructed in [B] a semi-stable genus 3 fibration over $\mathbf{P}^1$ with 5 singular fibres; each of these has 8 double points. Therefore

$$N = 40 \quad \text{and} \quad (4g+2)(s-2) = 42.$$

REFERENCES

- [B] A. BEAUVILLE: *Le nombre minimum de fibres singulières d’une courbe stable sur $\mathbf{P}^1$* . Astérisque **86** (1981), 97-108.
- [BKP] F. BOGOMOLOV, L. KATZARKOV, T. PANTEV: *Hyperelliptic Szpiro inequality*. Preprint math.GT/0106212.
- [S] L. SZPIRO: *Discriminant et conducteur des courbes elliptiques*. Séminaire sur les Pinceaux de Courbes Elliptiques (Paris, 1988). Astérisque **183** (1990), 7–18.
- [T] S.-L. TAN: *The minimal number of singular fibers of a semistable curve over $\mathbf{P}^1$* . J. Algebraic Geom. **4** (1995), 591–596.
- [X] G. XIAO: *Fibered algebraic surfaces with low slope*. Math. Ann. **276** (1987), 449–466.

Arnaud BEAUVILLE
Laboratoire J.-A. Dieudonné
UMR 6621 du CNRS
UNIVERSITÉ DE NICE
Parc Valrose
F-06108 NICE Cedex 02